

Refractive Surgery Overview



The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Refractive Surgery Overview



The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How bothersome would haloes around lights at night be? Because *some* aspect of the pt's post-op visual life will be less than ideal, key to successful refractive surgery is 1) developing a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are realistic and achievable.



Refractive Surgery Overview

The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How bothersome would haloes around lights at night be? Because *some* aspect of the pt's post-op visual life will be less than ideal, key to successful refractive surgery is 1) developing a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are *realistic and achievable*. Further, the surgeon must have a refined eye (ahem) for recognizing pre-existing ocular conditions that render the pt a poor surgical candidate (or increase the risk of complications down the road).

Refractive Surgery Overview



The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How bothersome would haloes around lights at night be? Because *some* aspect of the pt's post-op visual life will be less than ideal, key to successful refractive surgery is 1) developing a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are *realistic and achievable*. Further, the surgeon must have a refined eye (ahem) for recognizing pre-existing ocular conditions that render the pt a poor surgical candidate (or increase the risk of complications down the road).

Just as the goal of refractive surgery is simple, so too is its organization. Broadly speaking, there are but two types of surgery: **Corneal** and **intraocular**.

Refractive Surgery Overview



The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How bothersome would haloes around lights at night be? Because *some* aspect of the pt's post-op visual life will be less than ideal, key to successful refractive surgery is 1) developing a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are *realistic and achievable*. Further, the surgeon must have a refined eye (ahem) for recognizing pre-existing ocular conditions that render the pt a poor surgical candidate (or increase the risk of complications down the road).

Just as the goal of refractive surgery is simple, so too is its organization. Broadly speaking, there are but two types of surgery: **Corneal** and **intraocular**. Corneal (aka *keratorefractive*) procedures work by reshaping the cornea to offset the native refractive error of the eye; there are a plethora of techniques and technologies for doing this.

Refractive Surgery Overview



The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How bothersome would haloes around lights at night be? Because *some* aspect of the pt's post-op visual life will be less than ideal, key to successful refractive surgery is 1) developing a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are *realistic and achievable*. Further, the surgeon must have a refined eye (ahem) for recognizing pre-existing ocular conditions that render the pt a poor surgical candidate (or increase the risk of complications down the road).

Just as the goal of refractive surgery is simple, so too is its organization. Broadly speaking, there are but two types of surgery: **Corneal** and **intraocular**. Corneal (aka *keratorefractive*) procedures work by reshaping the cornea to offset the native refractive error of the eye; there are a plethora of techniques and technologies for doing this. *In contrast, intraocular procedures are more limited in their variety: They involve either replacing the native lens with an IOL, or placing in the anterior chamber an accessory IOL that offsets the eye's native refractive error.*




Refractive Surgery Overview

The goal of refractive surgery is deceptively straightforward and simple: To render the pt less reliant upon refractive accoutrements (ie, contacts and glasses). Ideally, a pt s/p refractive surgery would have 20/20 vision at *all* distances, under *any* lighting conditions, with *no* dysphotopsias (visual experiences that degrade vision quality), and with *no* risk of future negative repercussions vis a vis the long-term health and/or optical performance of the eye. Also ideally, the above could be achieved irrespective of pre-op refractive status and/or pre-existing ocular conditions.

Current technology is unable to meet this lofty ideal, and thus refractive surgery necessitates compromises and trade-offs; eg, If you *had* to pick one, would you rather be spectacle-free at distance, or near? Would it be acceptable if you only needed glasses in dimly-lit restaurants? How both *you* and *the pt* would be best served lightest might be? Proper assessment of the pt's post-op visual life *requires* a solid understanding of the pt's visual preferences and requirements, and 2) communicating effectively with the pt regarding what her post-op visual life will be; ie, establishing expectations that are realistic and achievable. Further, the surgeon must have a refined eye (ahem) for recognizing pre-existing ocular conditions that render the pt a poor surgical candidate (or increase the risk of complications down the road).


Let's start with a brief review of refractive error

Just as the goal of refractive surgery is simple, so too is its organization. Broadly speaking, there are but two types of surgery: **Corneal** and **intraocular**. Corneal (aka *keratorefractive*) procedures work by reshaping the cornea to offset the native refractive error of the eye; there are a plethora of techniques and technologies for doing this. In contrast, *intraocular procedures* are more limited in their variety: They involve either replacing the native lens with an IOL, or placing in the anterior chamber an accessory IOL that offsets the eye's native refractive error.




Refractive Surgery Overview

- The refractive state of an eye—that is, whether it is , or —is determined by the location of its two words




Refractive Surgery Overview

- The refractive state of an eye—that is, whether it is **emmetropic** , **myopic** or **hyperopic**—is determined by the location of its ***far point***



Refractive Surgery Overview


- The refractive state of an eye—that is, whether it is **emmetropic**, **myopic** or **hyperopic**—is determined by the location of its **far point**
- The far point is: *The location in space*
to the retina when the eye is not



Refractive Surgery Overview

- The refractive state of an eye—that is, whether it is **emmetropic**, **myopic** or **hyperopic**—is determined by the location of its **far point**
- The far point is: *The location in space conjugate to the retina when the eye is not accommodating*


(Accommodation refers to conformational changes in the ciliary body/lens to facilitate vision at near.)




Refractive Surgery Overview

- The refractive state of an eye—that is, whether it is **emmetropic**, **myopic** or **hyperopic**—is determined by the location of its **far point**
- The far point is: *The location in space conjugate to the retina when the eye is not accommodating*

(Accommodation refers to conformational changes in the ciliary body/lens to facilitate vision at near.)



Refractive Surgery Overview




(The Far Point of) **The Emmetropic Eye**




Refractive Surgery Overview

(The Far Point of) **The Emmetropic Eye**




In the **emmetropic** eye, the parallel rays from a location at infinity are focused to a point located precisely on the retina.




Refractive Surgery Overview

(The Far Point of) **The Emmetropic Eye**




In the **emmetropic** eye, the parallel rays from a location at infinity are focused to a point located precisely on the retina. *In other words, the far point of the emmetropic eye is at infinity.*

Far Point of the emmetropic eye: Infinity



Refractive Surgery Overview


(The Far Point of) **The Emmetropic Eye**




In the **emmetropic** eye, the parallel rays from a location at infinity are focused to a point located precisely on the retina. *In other words, the far point of the emmetropic eye is at infinity.* Thus, emmetropes see 20/20 (or better) at distance without correction.

Far Point of the emmetropic eye: Infinity

Refractive Surgery Overview




(The Far Point of) **The Myopic Eye**



Parallel rays
from infinity

In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.




Refractive Surgery Overview

(The Far Point of) The Myopic Eye




In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet .




Refractive Surgery Overview

(The Far Point of) The Myopic Eye




In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet **in the vitreous** .




Refractive Surgery Overview

(The Far Point of) The Myopic Eye




In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet **in the vitreous**. By the time they reach the retina, the rays have diverged to form a two words, not a focal point.



Refractive Surgery Overview


(The Far Point of) The Myopic Eye



Parallel rays
from infinity

In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet **in the vitreous** . By the time they reach the retina, the rays have diverged to form a **blur circle** , not a focal point.




Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length

much v
little




Parallel rays from infinity

In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet in the vitreous . By the time they reach the retina, the rays have diverged to form a blur circle , not a focal point.

You could say *the myopic eye has too much converging power for its length.*


much v
little



Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length



In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet in the vitreous. By the time they reach the retina, the rays have diverged to form a blur circle, not a focal point.


You could say *the myopic eye has too much converging power for its length*.



Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length




Note that if the retina was *here*, the rays from infinity would be focused to a point.

In contrast to the sharp uncorrected distance vision of the emmetrope, the blurred vision is the plight of the **myope**.

In the myopic eye, rays from infinity meet in the vitreous. By the time they reach the retina, the rays have diverged to form a blur circle, not a focal point.


You could say *the myopic eye has too much converging power for its length*.



Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length



In contrast to the sharp uncorrected distance vision of the emmetrope, consider the plight of the **myope**.

In the myopic eye, rays from infinity meet in the vitreous . By the time they reach the retina, the rays have diverged to form a blur circle , not a focal point.

You could say the myopic eye has too much converging power for its length.


To be focused on the retina, the Far Point of a myopic eye will have to offset its excess convergence with an equivalent amount of divergence. **To accomplish this...**




Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length




...the Far Point of a myopic eye is just anterior to the corneal plane.




Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length



...the **Far Point of a myopic eye is just anterior to the corneal plane.** Rays from this location are still quite divergent when they reach the eye, and this divergence offsets the excess convergence that is built into the myopic eye. Thus, rays originating from the far point end up sharply focused at the retina.




Refractive Surgery Overview

(The Far Point of) The Myopic Eye

The myopic eye has too much converging power for its length




...the Far Point of a myopic eye is just anterior to the corneal plane. Rays from this location are still quite divergent when they reach the eye, and this divergence offsets the excess convergence that is built into the myopic eye. Thus, rays originating from the far point end up sharply focused at the retina. This is why nearsighted individuals can read without glasses—they're able to put the material at or near their far point.




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye



Now consider the **hyperope**. Where do parallel rays meet in *these* eyes?


Refractive Surgery Overview



The Hyperopic Eye


Parallel rays from infinity (vergence = 0)

Parallel rays from infinity



This is where the rays would meet if they hadn't run into the retina.

Now consider the **hyperope**. Where do parallel rays meet in *these* eyes?
In the hyperopic eye, rays from infinity never meet—they run out of eyeball first.
Thus, like the myopic eye, the rays form a **blur circle**, not a focal point, at the retina.



Refractive Surgery Overview


(The Far Point of) The Hyperopic Eye

The hyperopic eye has too much v little converging power for its length

much v little

Parallel rays from infinity (vergence = 0)

Parallel rays from infinity



Now consider the **hyperope**. Where do parallel rays meet in *these* eyes? In the hyperopic eye, rays from infinity never meet—they run out of eyeball first. Thus, like the myopic eye, the rays form a blur circle, not a focal point, at the retina.

You could say **the hyperopic eye has too much v little converging power for its length.**


much v little



Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length



Now consider the **hyperope**. Where do parallel rays meet in *these* eyes?
In the hyperopic eye, rays from infinity never meet—they run out of eyeball first.
Thus, like the myopic eye, the rays form a blur circle, not a focal point, at the retina.
You could say **the hyperopic eye has too little converging power for its length.**




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length


Parallel rays from infinity (vergence = 0)

Parallel rays from infinity



Note that if the retina was *here*, the rays from infinity would be focused to a point.


Now consider the **hyperope**. Where do parallel rays meet? In the hyperopic eye, rays from infinity never meet—they form a blur circle. Thus, like the myopic eye, the rays form a blur circle, not a focal point, at the retina. You could say *the hyperopic eye has too little converging power for its length.*



Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length



Now consider the **hyperope**. Where do parallel rays meet in *these* eyes?
 In the hyperopic eye, rays from infinity never meet—they run out of eyeball first. Thus, like the myopic eye, the rays form a blur circle, not a focal point, at the retina. You could say the hyperopic eye has too *little* converging power for its length. In order to be conjugate to the retina, the Far Point of a hyperopic eye must contribute convergence to compensate for this lack of converging power.

To accomplish this...




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length

Far Point

Converging light



A and B are conjugate points


...the far point of a hyperopic eye is behind the corneal plane.



Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length



*...the far point of a hyperopic eye is **behind** the corneal plane. It contributes **convergence** to make up for the inadequate native convergence of the hyperopic eye.*




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length

Far Point

Converging light



A and B are conjugate points

...the far point of a hyperopic eye is *behind* the corneal plane. It contributes **convergence** to make up for the inadequate native convergence of the hyperopic eye. Thus, rays associated with the far point end up sharply focused at the retina.




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length

Converging light



A and B are conjugate points

...the far point of a hyperopic eye is *behind* the corneal plane. It contributes **convergence** to make up for the inadequate native convergence of the hyperopic eye. Thus, rays associated with the far point end up sharply focused at the retina. Don't get it twisted—hyperopes can't actually see behind their heads. (Do I really have to say that?)




Refractive Surgery Overview

(The Far Point of) The Hyperopic Eye

The hyperopic eye has too little converging power for its length


Converging light



...the far point of a hyperopic eye is *behind* the corneal plane. It contributes **convergence** to make up for the inadequate native convergence of the hyperopic eye. Thus, rays associated with the far point end up sharply focused at the retina. **Don't get it twisted—hyperopes can't actually see behind their heads. (Do I really have to say that?)** Unlike myopes—who can see at their far point just out in front of their faces—a hyperope is out of focus at **all** distances (absent correction or accommodation.)


Refractive Surgery Overview

- The myopic eye has too much power for its length, as we said




Myopic Eye






Refractive Surgery Overview

- The myopic eye has too much *converging* power for its length, as we said




Myopic Eye




Refractive Surgery Overview

- The myopic eye has too much *converging* power for its length, as we said




Think of it this way: The myopic eye refracts light as if an extra 'plus' lens was built into it. This so-called **error lens** contributes the excess convergence that produces a myopic refractive error.



Refractive Surgery Overview

- The myopic eye has too much **converging** power for its length, as we said




This explains why **myopes wear minus lenses** to correct their refractive error—minus lenses are needed to offset the excess convergence induced by the plus error lenses in their eyes.

Think of it this way: The **myopic eye refracts light as if an extra 'plus' lens was built into it.** This so-called **error lens** contributes the excess convergence that produces a myopic refractive error.

Refractive Surgery Overview


- The myopic eye has too much *converging* power for its length, as we said
- In contrast, the hyperopic eye has too much power for its length






Refractive Surgery Overview


- The myopic eye has too much *converging* power for its length, as we said
- In contrast, the hyperopic eye has too much *diverging* power for its length






Refractive Surgery Overview

- The myopic eye has too much *converging* power for its length, as we said
- In contrast, the hyperopic eye has too much *diverging* power for its length




Thus, the hyperopic eye acts as if it has a *minus* error lens within it, contributing the excess divergence resulting in a hyperopic refractive error.



Refractive Surgery Overview

- The myopic eye has too much *converging* power for its length, as we said
- In contrast, the hyperopic eye has too much *diverging* power for its length

This explains why **hyperopes wear plus lenses** to correct their refractive error—plus lenses are needed to offset the excess divergence induced by the minus error lenses in their eyes.





Thus, the **hyperopic eye acts as if it has a minus error lens within it**, contributing the excess divergence resulting in a hyperopic refractive error.




Refractive Surgery Overview

The goal of refractive surgery is to produce an error-lens offset that is incorporated into the eye itself, rather than worn on (CLs) or near (glasses) its anterior surface




Refractive Surgery Overview




As mentioned previously, refractive surgical procedures come in two basic forms—
 and


Refractive Surgery Overview



As mentioned previously, refractive surgical procedures come in two basic forms—*intraocular* and *corneal*.




Refractive Surgery Overview




Likewise, **intraocular procedures** come in two forms—, and

Refractive Surgery Overview




Likewise, **intraocular procedures** come in two forms—*pseudophakic* , and *phakic IOL* (PIOL)







Refractive Surgery Overview




A *pseudophakic procedure* involves removing the native lens and replacing it with an IOL powered to put parallel rays on the retina. The surgery itself is identical to that performed for cataracts. (Such procedures are referred to as three words .)








Refractive Surgery Overview




A *pseudophakic procedure* involves removing the native lens and replacing it with an IOL powered to put parallel rays on the retina. The surgery itself is identical to that performed for cataracts. (Such procedures are referred to as *clear lens extraction*.)







Refractive Surgery Overview



In a *phakic IOL procedure* the native lens is left in place, and a corrective lens is placed in front of it—an ‘*intraocular contact lens*’ if you will.




Refractive Surgery Overview





Phakic IOL

Refractive Surgery Overview




Phakic IOL


Refractive Surgery Overview



Phakic IOL vaulting over the native lens





Refractive Surgery Overview



Before we get into cornea-based refractive surgeries, let's take a look at [corneal optics](#)


Refractive Surgery Overview

The shape of the human cornea is *prolate*, meaning the central portion is steeper (ie, has a shorter radius of curvature) than the peripheral portion.





Refractive Surgery Overview

The shape of the human cornea is *prolate*, meaning the central portion is steeper (ie, has a shorter radius of curvature) than the peripheral portion.




Refractive Surgery Overview


The shape of the human cornea is *prolate*, meaning the central portion is steeper (ie, has a shorter radius of curvature) than the peripheral portion. On average, the central cornea is about 3-4D steeper than the periphery.



Refractive Surgery Overview





Power differential of central vs peripheral cornea
(don't memorize the numbers)



Refractive Surgery Overview


The anterior corneal surface adds about of convergence to incoming light (note how the air-cornea interface looks like a 'plus' lens)






Refractive Surgery Overview


The anterior corneal surface adds about **49D** of convergence to incoming light (note how the air-cornea interface looks like a 'plus' lens)






Refractive Surgery Overview






Refractive Surgery Overview






Refractive Surgery Overview



The net result across the cornea is an overall power of about +43D




Refractive Surgery Overview

The human eye averages about $+43\text{D}$ of total convergence, implying (correctly) that the cornea accounts for roughly $+49\text{D}$ of its focusing power




The net result across the cornea is an overall power of about $+43\text{D}$




Refractive Surgery Overview

The human eye averages about **60D** of total convergence, implying (correctly) that the cornea accounts for roughly **2/3** of its focusing power




The net result across the cornea is an overall power of about +43D




Refractive Surgery Overview

The human eye averages about **60D** of total convergence, implying (correctly) that the cornea accounts for roughly **2/3** of its focusing power




Of course, these are only averages. In order to perform keratorefractive surgery, one must have accurate measurements of central corneal power—ideally, at both its anterior and posterior surfaces.


The net result across the cornea is an overall power of about +43D



Refractive Surgery Overview


Two technologies are widely employed to determine central corneal power: Corneal , and corneal





Refractive Surgery Overview


Two technologies are widely employed to determine central corneal power: Corneal **topography** , and corneal **tomography**






Refractive Surgery Overview

Corneal topography works by reflecting a set of concentric rings (collectively called a two words) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings.




Refractive Surgery Overview

Corneal topography works by reflecting a set of concentric rings (collectively called a *Placido disk*) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings.



Refractive Surgery Overview


Corneal topography works by reflecting a set of concentric rings (collectively called a *Placido disk*) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings. Based on this analysis, the topographer creates a color-coded 'map' depicting the curvature across the corneal surface.



Refractive Surgery Overview


Corneal topography works by reflecting a set of concentric rings (collectively called a *Placido disk*) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings. Based on this analysis, the topographer creates a color-coded 'map' depicting the curvature across the corneal surface.





Refractive Surgery Overview

Corneal topography works by reflecting a set of concentric rings (collectively called a *Placido disk*) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings. Based on this analysis, the topographer creates a color-coded 'map' depicting the curvature across the corneal surface.




Corneal Placido-disk topography: Color map demonstrating *astigmatism* (ie, the cornea is steeper in its three meridian)



Refractive Surgery Overview

Corneal topography works by reflecting a set of concentric rings (collectively called a *Placido disk*) from the anterior corneal surface, and a computer analyzes the distances between, and shapes of, the reflected rings. Based on this analysis, the topographer creates a color-coded 'map' depicting the curvature across the corneal surface.




Corneal Placido-disk topography: Color map demonstrating *with-the-rule astigmatism* (ie, the cornea is steeper in its vertical meridian)

Refractive Surgery Overview

Corneal tomography works by mapping the anterior and posterior corneal surfaces in relation to one another. It allows for 3-D modeling of the cornea, including both anterior and posterior surface curvature and corneal thickness.








Refractive Surgery Overview

Corneal tomography works by mapping the anterior and posterior corneal surfaces in relation to one another. It allows for 3-D modeling of the cornea, including both anterior and posterior surface curvature and corneal thickness.

The data are acquired via one of two technologies (or both in combination):

-- **two-words**: A series of overlapping scans are directed at the cornea. The light reflects off both the anterior and posterior surfaces. These reflections are acquired and analyzed to produce a model of the central cornea.








Refractive Surgery Overview

Corneal tomography works by mapping the anterior and posterior corneal surfaces in relation to one another. It allows for 3-D modeling of the cornea, including both anterior and posterior surface curvature and corneal thickness.

The data are acquired via one of two technologies (or both in combination):

--*Scanning-slit* : A series of overlapping scans are directed at the cornea. The light reflects off both the anterior and posterior surfaces. These reflections are acquired and analyzed to produce a model of the central cornea.






Refractive Surgery Overview

Corneal tomography works by mapping the anterior and posterior corneal surfaces in relation to one another. It allows for 3-D modeling of the cornea, including both anterior and posterior surface curvature and corneal thickness.

The data are acquired via one of two technologies (or both in combination):

--*Scanning-slit*. A series of overlapping scans are directed at the cornea. The light reflects off both the anterior and posterior surfaces. These reflections are acquired and analyzed to produce a model of the central cornea.

- **eponym** *imaging*: A series of **eponym** images are taken and analyzed with respect to anterior and posterior corneal curvature and corneal thickness. The data from each image are knitted together to produce a model of the cornea.



Scheimpflug image of the cornea



Refractive Surgery Overview


Corneal tomography works by mapping the anterior and posterior corneal surfaces in relation to one another. It allows for 3-D modeling of the cornea, including both anterior and posterior surface curvature and corneal thickness.

The data are acquired via one of two technologies (or both in combination):

--*Scanning-slit*: A series of overlapping scans are directed at the cornea. The light reflects off both the anterior and posterior surfaces. These reflections are acquired and analyzed to produce a model of the central cornea.

--*Scheimpflug imaging*: A series of Scheimpflug images are taken and analyzed with respect to anterior and posterior corneal curvature and corneal thickness.


The data from each image are knitted together to produce a model of the cornea.



Scheimpflug image of the cornea




Refractive Surgery Overview



Pentacam corneal tomographer readout

Refractive Surgery Overview



Pentacam corneal tomographer readout


A) Anterior corneal values

- K_1 , K_2 , K_m : The two major meridians (K_1 , K_2). K_m is the average of K_1 and K_2
- R_f , R_s , R_m : Radii corresponding with K_1 , K_2 , and K_m , respectively
- QS: Quality score (I.e. "OK," "Data gaps," "Fix," "Model")
- Axis: The meridian that requires no cylinder power to correct astigmatism
- Astig: The central corneal astigmatism

B) Posterior corneal values


The same variables described for the back of the cornea.


C), D) Fuggedaboudit (too much for this overview)



Refractive Surgery Overview


In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a two words.





Refractive Surgery Overview


In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a *corneal ectasia* .






Refractive Surgery Overview


In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a *corneal ectasia*. An ectasia is a inflammatory vs non- condition characterized by progressive corneal bad change 1, the end result of which is corneal bad change 2. Pre-existing ectasia is a strong contraindication to many elective keratorefractive procedures, eg, LASIK.







Refractive Surgery Overview

In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a *corneal ectasia*. An ectasia is a noninflammatory condition characterized by progressive corneal thinning, the end result of which is corneal warpage. Pre-existing ectasia is a strong contraindication to many elective keratorefractive procedures, eg, LASIK.






Refractive Surgery Overview




In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a *corneal ectasia*. An ectasia is a noninflammatory condition characterized by progressive corneal thinning, the end result of which is corneal warpage. Pre-existing ectasia is a strong contraindication to many elective keratorefractive procedures, eg, LASIK. The two most common ectasias are and three words.





Refractive Surgery Overview

In addition to determining corneal power, pre-op corneal mapping is employed to determine whether a prospective keratorefractive pt has a *corneal ectasia*. An ectasia is a noninflammatory condition characterized by progressive corneal thinning, the end result of which is corneal warpage. Pre-existing ectasia is a strong contraindication to many elective keratorefractive procedures, eg, LASIK. The two most common ectasias are **keratoconus (KCN)** and **pellucid marginal degeneration (PMD)**.




Refractive Surgery Overview



KCN


Refractive Surgery Overview



Topography in KCN: Classic


three words

Refractive Surgery Overview



Topography in KCN: Classic *inferior corneal steepening*

Refractive Surgery Overview




Topography in PMD: Classic

two words

or


two diff words

Refractive Surgery Overview



Topography in PMD: Classic *kissing doves* or *crab claw*

Refractive Surgery Overview



Corneal Placido-disk topography: Mires typical of


KCN? PMD?

Refractive Surgery Overview




Corneal Placido-disk topography: Mires typical of KCN

Refractive Surgery Overview




Most *corneal refractive surgeries* involve altering the shape of the cornea in a way that impacts the vergence it imparts to incoming light.






Refractive Surgery Overview



Most *corneal refractive surgeries* involve altering the shape of the cornea in a way that impacts the vergence it imparts to incoming light. These alterations can involve [redacted] the cornea, [redacted] it, or some other means.


Refractive Surgery Overview



Most *corneal refractive surgeries* involve altering the shape of the cornea in a way that impacts the vergence it imparts to incoming light. These alterations can involve incising the cornea, lasering it, or some other means.




Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, LASIK), the cornea is reshaped so as to offset the effect of the error lens..



Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, LASIK), the cornea is reshaped so as to offset the effect of the error lens. In **myopic keratoablative surgery**, the central cornea is flattened v steepened to ↑ vs ↓ its converging power.






Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, LASIK), the cornea is reshaped so as to offset the effect of the error lens. In **myopic keratoablative surgery**, the central cornea is flattened to reduce its converging power.






Refractive Surgery Overview




Think of it as shaving down the peak of a mountain in order to make the structure more mesa-like

the cornea is reshaped so as to offset the surgery, **the central cornea is flattened**







Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, LASIK), the cornea is reshaped so as to offset the effect of the error lens. In **myopic keratoablative surgery**, the central cornea is flattened to reduce its converging power. **Hyperopic keratoablative surgery** is the opposite—the central cornea is flattened v steepened to ↑ vs ↓ its converging power.







Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, LASIK), the cornea is reshaped so as to offset the effect of the error lens. In **myopic keratoablative surgery**, the central cornea is flattened to reduce its converging power. **Hyperopic keratoablative surgery** is the opposite—the central cornea is *steepened* to *increase* its converging power.







Refractive Surgery Overview




In a *keratoablative laser procedures* (eg, L... effect of the error lens. In **myopic keratoa** to reduce its converging power. **Hyperop** the central cornea is **steepened** to incr

In contrast, hyperopic keratoablative surgery is akin to shaving down the rim of a mesa in order to make its structure more mountain-like








Refractive Surgery Overview



Note that, by definition, keratoablative refractive surgery involves reshaping the central cornea (and thereby altering its refractive power) via the removal (by annihilation) of corneal tissue





Refractive Surgery Overview

Refractive Surgery

Intraocular



Corneal

*One laser-based keratorefractive procedure does *not* involve tissue annihilation, rather, in it a section of corneal stroma is carved, then removed *en bloc*




Pseudophakic Phakic IOL Incisional **Laser** Other

Note that, by definition, keratoablative refractive surgery involves reshaping the central cornea (and thereby altering its refractive power) via **the removal (by annihilation) of corneal tissue***






Refractive Surgery Overview



What are the five laser-based keratorefractive procedures covered in the BCSC book?




Refractive Surgery Overview



What are the five laser-based keratorefractive procedures covered in the BCSC book?



Refractive Surgery Overview




What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago?



Refractive Surgery Overview




What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago? SMILE



Refractive Surgery Overview



In the SMILE procedure, the laser, with its ability to be focused at very precise depths within the cornea, is used to carve a segment (called a *lenticule*) of very specific shape within the stroma without disturbing the overlying or underlying tissue.


SMall-**I**ncision **L**enticule **E**xtraction — **SMILE**

What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago? **SMILE**



Refractive Surgery Overview



In the SMILE procedure, the femtosecond laser, with its ability to be focused at very precise depths within the cornea, is used to carve a segment (called a *lenticule*) of very specific shape within the stroma without disturbing the overlying or underlying tissue.


SMall-**I**ncision **L**enticule **E**xtraction — SMILE

What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago? **SMILE**



Refractive Surgery Overview




In the SMILE procedure, the femtosecond laser, with its ability to be focused at very precise depths within the cornea, is used to carve a segment (called a *lenticule*) of very specific shape within the stroma without disturbing the overlying or underlying tissue. The lenticule is then removed *en bloc* by being extracted through a very small incision (also created by the femto) that connects the femto-created intrastromal space and the corneal surface.

SMall-**I**ncision **L**enticule **E**xtraction — **SMILE**


What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago? **SMILE**

Refractive Surgery Overview



Creation of lenticule and small access (< 4 mm)



Removal of the lenticule




Refractive error is corrected

SMILE



Refractive Surgery Overview



In the SMILE procedure, the femtosecond laser, with its ability to be focused at very precise depths within the cornea, is used to carve a segment (called a *lenticule*) of very specific shape within the stroma without disturbing the overlying or underlying tissue. The lenticule is then removed *en bloc* by being extracted through a very small incision (also created by the femto) that connects the femto-created intrastromal space and the corneal surface. The resulting loss of tissue reshapes the central corneal surface in a way that produces a desired change in its refractive power.


SMall-**I**ncision **L**enticule **E**xtraction — **SMILE**

What are the five laser-based keratorefractive procedures covered in the BCSC book?

Which is the nonablative procedure referred to on a couple of slides ago? **SMILE**




Refractive Surgery Overview



In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an laser. But before the (same laser) can get to the stroma, the corneal epithelium has to get out of the way.




Refractive Surgery Overview



In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way.




Refractive Surgery Overview



In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. *The four keratoablative procedures differ solely in how the epithelium is handled.*



Refractive Surgery Overview




In **PRK**, the handling of the epithelium couldn't be more straightforward. It is simply cast aside—via scraping, chemical destruction, brushing, lasing, etc. This makes PRK the simplest of the laser keratorablative procedures: get the epithelium out of the way, then forget about it.

In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.



Refractive Surgery Overview




In **PRK**, the handling of the epithelium couldn't be more straightforward. It is simply cast aside—via scraping, chemical destruction, brushing, lasing, etc. This makes PRK the simplest of the laser keratorablative procedures: get the epithelium out of the way, then forget about it. However, PRK is associated with several post-operative complications that render it problematic, two of which are 1) it produces significant post-op [redacted], and 2) it is associated with an increased risk of post-op [redacted] —a potentially sight-threatening development.

In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)




Refractive Surgery Overview



In **PRK**, the handling of the epithelium couldn't be more straightforward. It is simply cast aside—via scraping, chemical destruction, brushing, lasing, etc. This makes PRK the simplest of the laser keratorablative procedures: get the epithelium out of the way, then forget about it. However, PRK is associated with several post-operative complications that render it problematic, two of which are 1) it produces significant post-op pain, and 2) it is associated with an increased risk of post-op haze formation—a potentially sight-threatening development.

In **keratoablative procedures**, remodeling of the central cornea occurs via annihilation of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)


Refractive Surgery Overview



Post-PRK haze



Refractive Surgery Overview




In contrast, the well-known **LASIK** procedure deals with the epithelium by doing an end-run around it.

In the case of LASIK, the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)



Refractive Surgery Overview




In contrast, the well-known **LASIK** procedure deals with the epithelium by doing an end-run around it. A two words is cut in the stroma and reflected, thereby moving the overlying epithelium out of the treatment area.

In of the the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)



Refractive Surgery Overview




In contrast, the well-known **LASIK** procedure deals with the epithelium by doing an end-run around it. A hinged flap is cut in the stroma and reflected, thereby moving the overlying epithelium out of the treatment area.

In of the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)



Refractive Surgery Overview



In contrast, the well-known **LASIK** procedure deals with the epithelium by doing an end-run around it. A hinged flap is cut in the stroma and reflected, thereby moving the overlying epithelium out of the treatment area. The underlying stromal bed is then lased, and the flap (with its intact epithelium) is laid back in place. Far less pain; vastly reduced risk of haze formation.

the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)

Refractive Surgery Overview



The diagram illustrates the four steps of LASIK surgery using cross-sectional views of an eye. Step 1 shows a microkeratome creating a flap. Step 2 shows the flap being lifted. Step 3 shows an excimer laser beam reshaping the cornea. Step 4 shows the flap being repositioned.

Step 1 : Corneal flap is created with a microkeratome.

Step 2 : The corneal flap is folded back.


Step 3 : Excimer laser beam reshapes the cornea.

Step 4 : The corneal flap is folded back in place.

LASIK



Refractive Surgery Overview




Like PRK, LASEK is a 'surface ablation' procedure. However, it deals very differently with the corneal epithelium.

In k
of t
na,
the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.



Refractive Surgery Overview




Like PRK, LASEK is a 'surface ablation' procedure. However, it deals very differently with the corneal epithelium. In LASEK, the epithelium is [redacted] and loosened by bathing it in [redacted].

In k
of t
a,
the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.



Refractive Surgery Overview




Like PRK, LASEK is a 'surface ablation' procedure. However, it deals very differently with the corneal epithelium. In LASEK, the epithelium is chemically devitalized and loosened by bathing it in alcohol.

In k
of t
na,
the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.





Refractive Surgery Overview



Like PRK, LASEK is a 'surface ablation' procedure. However, it deals very differently with the corneal epithelium. In LASEK, the epithelium is chemically devitalized and loosened by bathing it in alcohol. The loosened epithelium is then folded back, and the ablation is performed. Following the ablation, this 'epithelial flap' is smoothed back into place and covered with a bandage CL.

In k of t
 the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.


Refractive Surgery Overview



LASEK



Refractive Surgery Overview




Like PRK, LASEK is a 'surface ablation' procedure. However, it deals very differently with the corneal epithelium. In LASEK, the epithelium is chemically devitalized and loosened by bathing it in alcohol. The loosened epithelium is then folded back, and the ablation is performed. Following the ablation, this 'epithelial flap' is smoothed back into place and covered with a bandage CL. By re-positioning the epithelium, LASEK avoids the large epi defect (and resulting severe pain) of PRK.

In k
of t

the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.



Refractive Surgery Overview




Like LASEK, **epi-LASIK** is a surface-ablation variant designed to avoid the drawbacks of PRK. In it, a the first step in the procedure

In the first step in the procedure of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. [The four keratoablative procedures differ solely in how the epithelium is handled.](#)





Refractive Surgery Overview



Like LASEK, **epi-LASIK** is a surface-ablation variant designed to avoid the drawbacks of PRK. In it, a blunt keratome (an 'epikeratome') slides under the epithelium, separating it.

In [unclear] of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.


Refractive Surgery Overview



Epi-LASIK



Refractive Surgery Overview




Like LASEK, **epi-LASIK** is a surface-ablation variant designed to avoid the drawbacks of PRK. In it, a blunt keratome (an 'epikeratome') slides under the epithelium, separating it. The epithelial flap thus created is folded back, then re-placed after the stroma has been ablated. (BTW, *epipolis* is a Greek word meaning 'superficial'.)

In of the corneal stroma with an excimer laser. But before the excimer can get to the stroma, the corneal epithelium has to get out of the way. The four keratoablative procedures differ solely in how the epithelium is handled.



Refractive Surgery Overview


Refractive Surgery







Refractive Surgery Overview

Refractive Surgery






Refractive Surgery Overview




A set of incisions oriented radially (**radial keratotomy, RK**) will **flatten v steepen** the central cornea, and thereby **increase v reduce** its converging power. Thus, RK can be used to treat **myopia v hyperopia**.






Refractive Surgery Overview




A set of incisions oriented radially (**radial keratotomy, RK**) will flatten the central cornea, and thereby reduce its converging power. Thus, RK can be used to treat myopia .






Refractive Surgery Overview



Radial keratotomy




Refractive Surgery Overview



A set of incisions oriented radially (**radial keratotomy, RK**) will flatten the central cornea, and thereby reduce its converging power. Thus, RK can be used to treat myopia. Note that because incisions cannot produce overall *steepening* of the central cornea, RK cannot be used to treat hyperopia.




Refractive Surgery Overview



A set of incisions oriented radially (**radial keratotomy, RK**) will flatten the central cornea, and thereby reduce its converging power. Thus, RK can be used to treat myopia. Note that because incisions cannot produce overall *steepening* of the central cornea, RK cannot be used to treat hyperopia. Note also that, at this juncture, RK is considered obsolete, and is thus no longer performed in the US.




Refractive Surgery Overview




A set of incisions oriented radially (**radial keratotomy, RK**) will flatten the central cornea, and thereby reduce its converging power. Thus, RK can be used to treat myopia. Note that because incisions cannot produce overall *steepening* of the central cornea, RK cannot be used to treat hyperopia. Note also that, at this juncture, **RK is considered obsolete, and is thus no longer performed in the US.**

OTOH, *peripheral* incisional procedures to offset [redacted] refractive error are an important and oft-used surgical technique



Refractive Surgery Overview




A set of incisions oriented radially (**radial keratotomy, RK**) will flatten the central cornea, and thereby reduce its converging power. Thus, RK can be used to treat myopia. Note that because incisions cannot produce overall *steepening* of the central cornea, RK cannot be used to treat hyperopia. Note also that, at this juncture, **RK is considered obsolete, and is thus no longer performed in the US.**

OTOH, *peripheral* incisional procedures to offset *astigmatic* refractive error are an important and oft-used surgical technique



Refractive Surgery Overview




Specifically, this refers to **AK** and **LRI**

OTOH, *peripheral* incisional procedures to offset *astigmatic* refractive error are an important and oft-used surgical technique



Refractive Surgery Overview




Arcuate Keratotomy — AK
Limbal Relaxing Incisions — LRI

Both AK and LRI incisions are placed on the **steep v flat** meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the **paracentral v limbal** cornea, whereas LRI are made at the **paracentral K v limbal K**




Refractive Surgery Overview




Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies).

Refractive Surgery Overview




AK incisions


Limbal Relaxing Incisions



LR incisions



Refractive Surgery Overview




Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called **cross-linking**, the meridian 90 deg away.

flattening v steepening


flattening v steepening



Refractive Surgery Overview




Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called *coupling*, steepening the meridian 90 deg away.




Refractive Surgery Overview


Corneal Coupling Effect



Coupling




Refractive Surgery Overview



Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called *coupling*, steepening the meridian 90 deg away. The most common indication for AK is correction of astigmatism in an eye that is s/p surg procedure (two words).




Refractive Surgery Overview



Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called *coupling*, steepening the meridian 90 deg away. The most common indication for AK is correction of astigmatism in an eye that is s/p penetrating keratoplasty .




Refractive Surgery Overview



Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called *coupling*, steepening the meridian 90 deg away. The most common indication for AK is correction of astigmatism in an eye that is s/p penetrating keratoplasty. LRI are typically placed at the time of surgery to correct corneal astigmatism in hopes of producing better UCVA post-op.



Refractive Surgery Overview




Both AK and LRI incisions are placed on the steep meridian of the cornea, in pairs located on opposite sides of the cornea. AK incisions are made in the paracentral cornea, whereas LRI are made at the limbus (as their name implies). Both techniques have the effect of flattening the meridian in which they're placed, but through a process called *coupling*, steepening the meridian 90 deg away. The most common indication for AK is correction of astigmatism in an eye that is s/p penetrating keratoplasty. LRI are typically placed at the time of cataract surgery to correct corneal astigmatism in hopes of producing better UCVA post-op.



Refractive Surgery Overview


Refractive Surgery





Refractive Surgery Overview

Refractive Surgery




These are the 'other' procedures covered in the *BCSC* book

- Conductive Keratoplasty — **CK**
- Small Aperture Inlay — **SAI**
- Corneal Reshaping Inlay — **CRI**
- Corneal **CROSS** Linking — **CXL**
- Intrastromal Corneal Ring Segments — **ICRS**



Refractive Surgery Overview

Refractive Surgery



These three treat




- Conductive Keratoplasty — CK
- Small Aperture Inlay — SAI
- Corneal Reshaping Inlay — CRI
- Corneal CROSS Linking — CXL
- Intrastromal Corneal Ring Segments — ICRS




Refractive Surgery Overview

Refractive Surgery




These three treat presbyopia



- Conductive Keratoplasty — **CK**
- Small Aperture Inlay — **SAI**
- Corneal Reshaping Inlay — **CRI**
- Corneal CROSS Linking — **CXL**
- Intrastromal Corneal Ring Segments — **ICRS**




Refractive Surgery Overview





Refractive Surgery Overview

Refractive Surgery





Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal **flattening v steepening** thereby increasing **convergence v divergence** and improving vision at near.

Conductive Keratoplasty — **CK**

Small Aperture Inlay — **SAI**

Corneal Reshaping Inlay — **CRI**

Corneal **CROSS** Linking — **CXL**

Intrastromal **Corneal Ring Segments** — **ICRS**



Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal steepening, thereby increasing convergence and improving vision at near.

Conductive Keratoplasty — **CK**

Small Aperture Inlay — SAI

Corneal Reshaping Inlay — CRI

Corneal **CROSS** Linking — CXL

Intrastromal **Corneal Ring Segments** — ICRS


Overview



CK probe



CK in action



CK scars



Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal steepening, thereby increasing convergence and improving vision at near.

The **SAI** is a [describe it—size, shape] that is implanted in [where]. Its central [] produces a [] effect to improve near vision.

- Conductive Keratoplasty — CK
- Small Aperture Inlay — SAI**
- Corneal Reshaping Inlay — CRI
- Corneal **CROSS** Linking — CXL
- Intrastromal **Corneal Ring Segments** — ICRS



Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal steepening, thereby increasing convergence and improving vision at near.

The **SAI** is a tiny donut-shaped disc that is implanted in the central corneal stroma. Its central aperture produces a pinhole effect to improve near vision.

- Conductive Keratoplasty — **CK**
- Small Aperture Inlay — SAI**
- Corneal Reshaping Inlay — **CRI**
- Corneal **CROSS** Linking — **CXL**
- Intrastromal **Corneal Ring Segments** — **ICRS**



Refractive Surgery Overview




Made from Polyvinylidene Fluoride (PVDF)




SAI

Refractive Surgery Overview



SAI



Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal steepening, thereby increasing convergence and improving vision at near.

The **SAI** is a tiny donut-shaped disc that is implanted in the central corneal stroma. Its central aperture produces a pinhole effect to improve near vision.

Like the SAI, the **CRI** is a tiny device implanted in the central corneal stroma. However, its mechanism of action is different—it is **shape** with a central **shape** that **increases v decreases** the curvature of the overlying corneal surface, which in turn increases corneal **convergence v divergence** and thus improves near vision.

- Conductive Keratoplasty — CK
- Small Aperture Inlay — SAI
- Corneal Reshaping Inlay — CRI**
- Corneal CROSS Linking — CXL
- Intrastromal Corneal Ring Segments — ICRS



Refractive Surgery Overview

Refractive Surgery

Intraocular

Corneal

Pseudophakic

In **CK**, a thermal probe is used to produce a set of focal corneal scars. The scars produce local corneal steepening, thereby increasing convergence and improving vision at near.

The **SAI** is a tiny donut-shaped disc that is implanted in the central corneal stroma. Its central aperture produces a pinhole effect to improve near vision.

Like the SAI, the **CRI** is a tiny device implanted in the central corneal stroma. However, its mechanism of action is different—it is disc-shaped with a central 'bump' that increases the curvature of the overlying corneal surface, which in turn increases corneal convergence and thus improves near vision.

Conductive Keratoplasty — **CK**



Small Aperture Inlay — **SAI**

Corneal Reshaping Inlay — **CRI**

Corneal CROSS Linking — **CXL**

Intrastromal Corneal Ring Segments — **ICRS**


Refractive Surgery Overview



CRI



Refractive Surgery Overview




CXL involves the induction of a chemical reaction that [redacted]. The result is a stabilization of the keratoconus process.

Corneal CROSS Linking — CXL

Intrastromal Corneal Ring Segments — ICRS



Refractive Surgery Overview




CXL involves the induction of a chemical reaction that strengthens the bonds between corneal stroma fibrils . The result is a stabilization of the keratoconus process.

Corneal CROSS Linking — CXL




Refractive Surgery Overview

**BEFORE CXL : LESS CROSSLINKING
= WEAKER CORNEA**




**AFTER CXL : MORE CROSSLINKING
= STRONGER CORNEA**




CXL concept




Refractive Surgery Overview




1. We remove the Epithelium



2. Ribloflavin (Vitamin B2) eye drops are applied onto the cornea



3. 1 minute later, the solution is irrigated or washed away by the surgeon



4. An ultra-violet light (UVA) illuminates the Riboflavin solution for the corneal cross-linking procedure

CXL: Process


Refractive Surgery Overview



CXL: Process



Refractive Surgery Overview



CXL involves the induction of a chemical reaction that strengthens the bonds between corneal stroma fibrils. The result is a stabilization of the keratoconus process.


ICRS employs shape segments of abb.. These segments are placed in the peripheral corneal layer, where they produce local flattening v steepening

CK
SAI
CRI
CXL

Intrastromal Corneal Ring Segments — **ICRS**




Refractive Surgery Overview



CXL involves the induction of a chemical reaction that strengthens the bonds between corneal stroma fibrils . The result is a stabilization of the keratoconus process.
ICRS employs semicircular segments of PMMA . These segments are placed in the peripheral corneal stroma , where they produce local flattening .



Intrastromal Corneal Ring Segments — **ICRS**

Refractive Surgery Overview





Intrastromal ring segments

Refractive Surgery Overview



Intrastromal ring segments *in situ*

Refractive Surgery Overview



Intrastromal ring segments placed for KCN